




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 1/00, 19/06, 19/16		A1	(11) International Publication Number: WO 95/32980
			(43) International Publication Date: 7 December 1995 (07.12.95)
(21) International Application Number: PCT/US95/06825		(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).	
(22) International Filing Date: 26 May 1995 (26.05.95)			
(30) Priority Data: 08/249,442 26 May 1994 (26.05.94) US			
(60) Parent Application or Grant (63) Related by Continuation US 08/249,442 (CIP) Filed on 26 May 1994 (26.05.94)		Published With international search report.	
(71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): RAVIKUMAR, Vasulinga [IN/US]; 2325 Via Franciscas #J, Carlsbad, CA 92008 (US). COLE, Douglas, L. [US/US]; 4992 Smith Canyon Court, San Diego, CA 92130 (US).			
(74) Agent: CALDWELL, John, W.; Woodcock Washburn Kurtz Mackiewicz & Norris, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).			

(54) Title: SYNTHESIS OF OLIGONUCLEOTIDES

(57) Abstract

Synthetic processes are provided for the solution phase synthesis of oligonucleotides, especially phosphorothioate oligonucleotides, and intermediate compounds useful in the processes. Intermediates having structure (I) are prepared in accordance with preferred embodiments.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

SYNTHESIS OF OLIGONUCLEOTIDES

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation in part of United States Serial No. 08/099,075 filed July 29, 1993, and United States Serial No. 08/249,442 filed May 26, 1994 each assigned to the assignee of this invention.

FIELD OF THE INVENTION

This invention is directed to processes for synthesizing oligonucleotides, especially phosphorothioate oligonucleotides, and to intermediates used in that process. This invention is drawn to solution phase syntheses having improved efficiencies and enhanced convenience and cost.

BACKGROUND OF THE INVENTION

Oligonucleotides are important materials for research, diagnostic, therapeutic and other purposes. An ever-growing demand for improved oligonucleotides, oligonucleotide analogs and for methods for their preparation and use has arisen. For example, oligonucleotides are widely used in genomic research as probes, primers and a wide array of other research uses. One widely used technique that uses oligonucleotides primers is PCR (polymerase chain reaction) amplification.

Oligonucleotides are also useful in diagnostics since they can specifically hybridize to nucleic acids of interest in the etiology of disease. Oligonucleotides are currently in clinical trials as therapeutic moieties in the

- 2 -

treatment of disease states. For example, workers in the field have now identified oligonucleotide compositions that are capable of modulating expression of genes implicated in viral, fungal and metabolic diseases. In short,

5 oligonucleotides are important molecules having a large commercial impact in biotechnology and medicine. Improved methods for the synthesis of oligonucleotides are in demand, especially methods which have improvements in cost and convenience.

10 The current methods of choice for the preparation of phosphorothioate oligonucleotides employ solid-phase synthesis wherein an oligonucleotide is prepared on a polymer or other solid support. Solid-phase synthesis relies on sequential addition of nucleotides to one end of a
15 growing oligonucleotide. Typically, a first nucleoside is attached to an appropriate support, e.g. glass, and nucleotide precursors, typically phosphoramidites, are added stepwise to elongate the growing oligonucleotide. The nucleotide phosphoramidites are conventionally reacted with
20 the growing oligonucleotide using the principles of a "fluidized bed" for mixing of the reagents. The silica supports suitable for anchoring the oligonucleotide are very fragile and thus can not be exposed to aggressive mixing.

In these and other solid-phase procedures the
25 oligonucleotide is synthesized as an elongating strand. However, the number of individual strands that can be anchored to a unit surface area of the support is limited. Also, the commercially available activated nucleotides that are presently used to add to a growing oligonucleotide are
30 relatively expensive and must be used in stoichiometric excess. Also, the activating agents, e.g. tetrazole, are used in large excess.

The chemical literature discloses numerous processes for coupling nucleosides through phosphorous-
35 containing covalent linkages to produce oligonucleotides of defined sequence. One of the most popular processes is the phosphoramidite technique (see, e.g., Beaucage, et al.,

- 3 -

Tetrahedron 1992, 48, 2223 and references cited therein), wherein a nucleoside or oligonucleotide having a free hydroxyl group is reacted with a protected cyanoethyl phosphoramidite monomer in the presence of a weak acid to
5 form a phosphite-linked structure. Oxidation of the phosphite linkage followed by hydrolysis of the cyanoethyl group yields the desired phosphodiester or phosphorothioate linkage.

The phosphoramidite technique, however, is not
10 without its disadvantages. For example, cyanoethyl phosphoramidite monomer is quite expensive. Although considerable quantities of monomer go unreacted in a typical phosphoramidite coupling, unreacted monomer can be recovered, if at all, only with great difficulty. Also,
15 acrylonitrile, the by-product of deprotection of the cyanoethoxy group on the phosphate group is carcinogenic and in some cases acts as a Michael acceptor to form undesired side-products.

Other exemplary solid state synthetic schemes are
20 set forth in U.S. Patents RE. 34,069 Koster et al.; and 5,132,418 Carruthers et al..

While currently utilized solid phase syntheses are useful for preparing small quantities of oligonucleotide, they typically are not amenable to the preparation of large
25 quantities of oligonucleotides necessary for biophysical studies, pre-clinical and clinical trials and commercial production. Moreover, such synthetic procedures are very expensive. Thus, although there is a great demand for oligonucleotides, especially phosphorothioate
30 oligonucleotides, the art suggests no large scale techniques for their preparation. Accordingly, there remains a long-felt need for such methods and for intermediates useful in such methods.

- 4 -

BRIEF DESCRIPTION OF THE DRAWINGS


Figure 1 is a reaction schematic depicting exemplary routes for the syntheses of intermediates useful in the practice of the invention.

5 Figure 2 is a schematic showing exemplary coupling of the intermediates of Figure 1.


Figure 3 depicts the coupling of the moieties of Figure 2.

SUMMARY OF THE INVENTION

10 In accordance with the invention, methods are provided for the synthesis of oligonucleotide moieties. A first synthon, having the structure




is caused to react with a second synthon having the structure



15 to form a moiety having the structure

- 5 -



Wherein each Q is independently O, S, CH₂, CHF or CF₂. In accordance with embodiments of the invention, each B_x is independently a nucleosidic base; each X is independently, OH, SH, SCH₃, F, OCN, O(CH₂)_nNH₂, O(CH₂)_nCH₃, where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl or aralkyl; Cl, Br, CN, CF₃, OCF₃, O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide; Y is a 5' hydroxyl protecting group; W is a 3' hydroxyl protecting group; each Z is O or S; T is a phosphorous blocking group; U is a phosphite activating group; and n is an integer from 0 to 50.

In accordance with preferred embodiments, each group T can be a silylalkoxy group wherein the Si atom of this group includes three substituent R groups thereon. Preferably these R groups, independently, are alkyl or aryl with methyl, t-butyl and phenyl being most preferred. U can

- 6 -

be a dialkylamino group. In another preferred embodiment, the first synthon is formed by coupling of a 3' hydroxyl protected nucleoside with a 5' protected nucleoside through a phosphite linkage.

5 In yet other preferred embodiments, the processes of the invention are accomplished such that the second synthon is prepared via reaction of a precursor nucleoside with a reagent $[R_1R_2N]_2PO(CH_2)_xSiR_3R_4R_5$ wherein R_1 and R_2 independently are alkyl having 1 to about 10 carbon atoms,
10 R_3 , R_4 , and R_5 are, independently, alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbons atoms, and x is 1 to about 7. Useful as the precursor nucleoside for the preparation of such a second synthon is a nucleoside that corresponds in structure to the second synthon above,
15 except it bears a hydrogen atom on its 3' hydroxyl group in place of the U-P-T groups. Preferably this reaction is conducted in the presence of 1H-Tetrazole, 5-(4-Nitrophenyl)-1H-tetrazole, or diisopropylammonium tetrazolide.


20 It is also preferred that the processes further comprise removing the groups W, T, and Y from the moiety and oxidizing the moiety to form either phosphorothioate or phosphodiester bonds.

The processes of the invention may be performed
25 iteratively such that the product resulting from a first reaction sequence is transformed into a "new" first synthon for iterative reaction with a further second synthon by removal of the Y group. The processes of the invention may further be performed iteratively such that the product
30 resulting from a first reaction sequence is transformed into a "new" second synthon for iterative reaction with a further first synthon by removal of the W group followed by phosphitation. While solution phase reactions are preferred in the practice of this invention, in iterative processes
35 and otherwise, one of the "new" second synthons of the above process, as described in the preceding sentence, can be used in the place of a standard mono-nucleotide phosphoramidite

- 7 -

synthon in a standard solid phase reaction to elongated an growing oligonucleotide on a solid phase support. Thus instead of adding just a single nucleotide at a time as is practiced in current solid phase synthesis technology, 5 dimers, trimers, tetramers and even high homologues can be advantageously linked to a growing oligonucleotide on a solid support. Such dimers, trimers, tetramers and high homologues are prepared using the processes of the invention and then are carried over and used in place of the standard 10 phosphoamidites of currently practiced solid phase oligonucleotide synthesis.


In accordance with other embodiments of the invention, there are reacted together, in solution, a first synthon comprising at least two nucleoside units having a 5' 15 location protected with a 5' hydroxylic blocking group and a 3' location substituted with a function having the formula




wherein U is a phosphite activating group and T is a phosphorous blocking group, with a second synthon comprising 20 a nucleoside unit having a 3' location protected with a 3' hydroxylic blocking group and a 5' location capable of reacting with the U-P-T function. It is preferred that the product of the reaction be oxidized to form either phosphodiester or phosphorothioate internucleoside bonds. 25 It is also preferred for some embodiments that the second synthon comprise at least two nucleoside units.

In other preferred methodologies, the function U-P-T is incorporated through reaction with a reagent $[\text{R}_1\text{R}_2\text{N}]_2\text{PO}(\text{CH}_2)_x\text{SiR}_3\text{R}_4\text{R}_5$ wherein R_1 and R_2 independently are 30 alkyl having 1 to about 10 carbon atoms, R_3 , R_4 , and R_5 are, independently, alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbons atoms, and x is 1 to about 7.

In other aspects of the invention, compounds are 35 provided having the formulas



and



Where each Q is, independently O, S, CH₂, CHF or CF₂. In such compounds, each B_x is independently a nucleosidic base; each X is independently, OH, SH, SCH₃, F, OCN, O(CH₂)_nNH₂, O(CH₂)_nCH₃ where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl or aralkyl; Cl, Br, CN, CF₃, OCF₃, O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide; each Z is independently O or S; Y is H or a hydroxyl protecting group; U is a phosphite activating group; each T is independently a phosphorous blocking group; and n is an integer from 1 to 50.

20 In accordance with other embodiments of the invention, the phosphorous blocking group has the formula - $O(CH_2)_xSiR_3R_4R_5$ wherein R_3 , R_4 , and R_5 are, independently, alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbons atoms, and x is 1 to about 7.


25 It is a significant aspect of the present invention that libraries of oligomers can be constructed and stored to facilitate the synthesis and identification of useful oligonucleotides. Accordingly, the invention is

- 9 -

directed to libraries comprising a plurality of compounds in accordance with the foregoing formulas and to oligonucleotides synthesized therefrom.

In a further aspect of the invention methods are provided for the preparation of a plurality of target oligonucleotides, each target oligonucleotide comprising at least one common nucleotide subsequence, comprising:

identifying said common nucleotide subsequences;
providing an oligonucleotide synthon of formula:



wherein:

each Q is O;
each B_x is independently a nucleosidic base;
15 each X is independently, H, OH, F, O-alkyl or substituted O-alkyl wherein said substitution is amino, alkoxy or nitrogen heterocycle;
each Z is independently O or S;
Y is H or a hydroxyl protecting group;
20 U is a phosphite activating group;
each T is independently a phosphorous blocking group; and

n is an integer from 1 to 200;
said oligonucleotide synthon having one of said common
25 nucleotide subsequences; and

reacting said synthon with nucleoside precursors to form said target oligonucleotide. Preferably, the reacting is performed in an organic solvent, preferably

- 10 -


below the boiling point of the solvent. In some preferred embodiments a plurality of said synthons are so reacted.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS


This invention provides new and improved processes
5 for the preparation of phosphorous-containing covalent linkages and intermediates useful in such processes. Utilizing these processes and intermediates, phosphodiester and phosphorothioate oligonucleotides are prepared from a plurality of nucleoside or oligonucleotide subunits. The
10 nucleoside subunits may be "natural" or "synthetic" moieties. Thus, in the context of this invention, the term "oligonucleotide" in a first instance refers to a polynucleotide formed from a plurality of linked nucleoside units. The nucleosides are formed from naturally occurring
15 bases and pentofuranosyl sugar groups. The term "oligonucleotide" thus effectively includes naturally occurring species or synthetic species formed from naturally occurring subunits.

The present invention provides processes for the
20 efficient synthesis of the oligonucleotides and modified oligonucleotide compounds (analogues). In preferred embodiments, certain products of the invention are prepared by processes that comprise contacting a protected nucleoside having formula I, i.e. a precursor nucleoside, with a
25 phosphitylation reagent having the formula $(R_1R_2N)_2PO(CH_2)_xSiR_3R_4R_5$ for a time and under reaction conditions effective to form a silylalkyl phosphoramidite monomer having formula II

- 11 -



I



II

wherein:

- 5 each Q is, independently, O, S, CH₂, CHF or CF₂;
 each B_x is, independently, a nucleosidic base;
 each X is, independently, OH, SH, SCH₃, F, OCN,
 O(CH₂)_nNH₂, O(CH₂)_nCH₃ where n is from 1 to about 10; C₁ to C₁₀
 lower alkyl, substituted lower alkyl, alkaryl or aralkyl;
 10 Cl, Br, CN, CF₃, OCF₃, O-, S-, or N-alkyl; O-, S-, or N-
 alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl;
 heterocycloalkaryl; aminoalkylamino; polyalkylamino; sub-
 stituted silyl; an RNA cleaving group; a conjugate; a
 15 reporter group; an intercalator; a group for improving the
 pharmacokinetic properties of an oligonucleotide; or a group
 for improving the pharmacodynamic properties of an
 oligonucleotide;

Y is a 5' hydroxyl protecting group;


- 20 R₁ and R₂, independently, are alkyl having 1 to
 about 10 carbon atoms;

R₃, R₄, and R₅ are, independently, alkyl having 1 to
 about 10 carbon atoms or aryl having 6 to about 10 carbons
 atoms; and


x is 1 to about 7.

- 25 Monomers of formula II then are contacted with a
 3' hydroxyl protected nucleoside having formula III for a
 time and under reaction conditions effective to form dimers
 having formula IV

- 12 -



III




IV


5 where W is a 3' hydroxyl protecting group and Y is a 5' hydroxyl protecting group and R_3 , R_4 and R_5 are defined as above.

A dimer having formula IV is then contacted with an oxidizing agent for a time and under reaction conditions effective to form oxidation products having formula V (where Z = O or S and $n = 1$). The dimer of formula V can be converted into a dimeric synthon analogous to either the monomeric synthon of structure II or to the monomeric synthon of structure III. Alternatively subsequently
 10 reaction of the dimeric compound of structure V with fluoride ion in the conventional manner gives either a phosphodieser or phosphorothioate moiety having formula VI.

- 13 -




V




VI

For conversion of the dimer of formula V (where n is 1) to a dimeric synthon analogous to that of structure III above, the 5' hydroxy blocking group Y is removed in a conventional manner to yield a compound of formula VII. This compound is then subsequently contacted with a further phosphitylated nucleoside of formula II under reaction conditions effective to form trimers of formula VIII.



VII




VIII


- 14 -

Trimers having formula VIII are then contacted with an oxidizing agent for a time and under reaction conditions effective to form oxidation products having formula V (where n is increased by one). If a trimeric product is desired, the trimer of formula V is deblocked such as with the conventional fluoride ion or by the use of tetrafluorosilane in a "wet" solvent to produce either a phosphodiester or phosphorothioate moiety having formula VI. If higher homologues are desired, the trimer of formula VIII can be deblocked at the 5' position removing the Y hydroxyl protecting group to yield a synthon analogous to formula VII or deblocked at the 3' position removing the W hydroxyl protecting group and phosphitylated to yield a trimeric synthon analogous to the compound of formula II. Thus a library of building blocks of various lengths can be formed. Units can be added together as monomers, as dimers, as trimers, etc, for the preparation of higher homologues.

In other embodiments of the present invention, compounds of formula I are contacted with other suitable coupling reagents to form monomers of formula IIa, wherein U is a phosphite activating group, and T is a phosphorous blocking group. Monomers of formula IIa then are contacted with 3' hydroxyl protected nucleoside having formula III for a time and under reaction conditions effective to form dimers having formula IVa.



IIa




IVa



- 15 -

where W is a 3' hydroxyl protecting group, Y is a 5' hydroxyl protecting group and T is a phosphorous protecting group.

A dimer having formula IVa is then contacted with an oxidizing agent for a time and under reaction conditions effective to form oxidation products having formula Va (where Z = O or S and n = 1).



A dimer of formula Va can also be reacted conventionally to remove 5' hydroxy blocking group Y to yield a compound of formula VIIa (where n is 1) and subsequently contacted with a further phosphitylated nucleoside of formula II under reaction conditions effective to form trimers of formula VIIIa.





5

Trimers having formula VIIIa are then contacted with an oxidizing agent for a time and under reaction conditions effective to form oxidation products having formula Va (where n is increased by one). This cycle is repeated to give the length of the desired product. If such product is the final product it is subsequently reacted such as with the conventional fluoride ion or tetrafluorosilane in a "wet" solvent to produce either a phosphodiester or phosphorothioate moiety having formula VI.

15 Oligonucleotides of formula Va can also be converted to higher analogues corresponding to phosphitylated nucleosides of formula II, above. To form such compounds a dimer of formula Va (n = 1) is deprotected at its 3' hydroxyl by removal of the group W and
20 phosphitylated to yield a dimer of formula Vb. The compound of formula Vb can be reacted with a compound of formula VIIa to form a new compound Vc, having one phosphite linking group located between the portion obtained from the compound

of formula Vb and the portion obtained from the compound of formula VIIa.



Subsequent oxidization of the phosphite linkage in Vc to a phosphate linkages yields a new compound of formula Va where n is the total of the n units from both the compound of formula VIIa and the compound of formula Vb.

10 Oligonucleotides according to the invention also
can include modified subunits. Representative modifications
include modification of a heterocyclic base portion of a
nucleoside or a sugar portion of a nucleoside. Exemplary
modifications are disclosed in the following U.S. Patent
15 Application: Serial No. 463,358, filed January 11, 1990,
entitled "Compositions And Methods For Detecting And
Modulating RNA Activity; Serial No. 566,977, filed August
13, 1990, entitled "Sugar Modified Oligonucleotides That

- 18 -

Detect And Modulate Gene Expression; Serial No. 558,663, filed July 27, 1990, entitled "Novel Polyamine Conjugated Oligonucleotides"; Serial No. 558,806, filed July 27, 1991, entitled "Nuclease Resistant Pyrimidine Modified
5 Oligonucleotides That Detect And Modulate Gene Expression"; and Serial No. PCT/US91/00243, filed January 11, 1991, entitled "Compositions and Methods For Detecting And Modulating RNA Activity". Each of these patent applications are assigned to the assignee of this invention. The
10 disclosure of each is incorporated herein by reference.

The term oligonucleotide thus refers to structures that include modified portions, be they modified sugar moieties or modified base moieties, that function similarly to natural bases and natural sugars. Representative
15 modified bases include deaza or aza purines and pyrimidine used in place of natural purine and pyrimidine bases; pyrimidine having substituent groups at the 5 or 6 position; purines having altered or replacement substituent groups at the 2, 6, or 8 positions. Representative modified sugars
20 include carbocyclic or acyclic sugars, sugars having substituent groups at their 2' position, and sugars having substituents in place of one or more hydrogen atoms of the sugar. Other altered base moieties and altered sugar moieties are disclosed in U.S. Patent No. 3,687,808 and PCT
25 application PCT/US89/02323.

In certain embodiments, the compounds of the invention include conjugate groups. Conjugate groups of the invention include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups
30 that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugate groups include cholesterols, phospholipids, biotin, phenanthroline, phenazine, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines,
35 coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this invention, include groups that improve oligomer uptake, enhance

- 19 -

oligomer resistance to degradation, and/or strengthen sequence-specific hybridization with RNA. Groups that enhance the pharmacokinetic properties, in the context of this invention, include groups that improve oligomer uptake, distribution, metabolism or excretion. Representative conjugate groups are disclosed in International Patent Application PCT/US92/09196, filed October 23, 1992, United States Patent Application Serial No. 116,801, filed September 3, 1993, and United States Patent No. 5,218,105. Each of the foregoing is commonly assigned with this application. The entire disclosure of each is incorporated herein by reference.

Altered base moieties or altered sugar moieties also include other modifications consistent with the spirit of this invention. Such oligonucleosides are best described as moieties that are structurally distinguishable from yet functionally interchangeable with naturally occurring or synthetic wild type oligonucleotides. All such oligonucleotides are comprehended by this invention so long as they function effectively to mimic the structure of a desired RNA or DNA strand.

Oligonucleotides prepared via the process and intermediates of the invention can be used as primers for enzymes which polymerize nucleotides, for example, polymerases and reverse transcriptases, such as in the well-known PCR technology. Oligonucleotides can also be used as probes for specific oligonucleotide sequences. Conventional methods for use of oligonucleotides as probes are disclosed in, for example, "Molecular Cloning; A Laboratory Manual" 2d. Ed., J. Sambrook et al., Cold Spring Harbor Press, at chapter 10 and the references therein, which are hereby incorporated by reference.

While compounds of the invention as small as dimeric units are useful, preferred the oligonucleotides of the invention comprise from about 10 to about 30 subunits, although oligomers of up to 50 or even several hundred subunits may be useful. It is more preferred that such

- 20 -

oligonucleotides comprise from about 15 to 25 subunits. As will be appreciated, a subunit is a base and sugar combination suitably bound to adjacent subunits through a phosphorous-containing linkage. When used as "building blocks" in assembling oligonucleotides, even smaller assemblies are preferred. The smallest, a dinucleotide assembly, is two nucleotide linked by a protected phosphorothioate or phosphodiester linkage.

One use of oligonucleotides is in targeting RNA or DNA. One specific such use is targeting of RNA or DNA that can code for protein. It is preferred that the RNA or DNA portion which is targeted using the oligonucleotide be preselected to comprise that portion of DNA or RNA which codes for the protein whose formation or activity is to be modulated. The targeting portion of the composition to be employed is, thus, selected to be complementary to the preselected portion of DNA or RNA. Such use that is to be an antisense oligonucleotide for that portion.

In accordance with one preferred embodiment of this invention, a compound described as Oligonucleotide 5320 having the structure $T_2G_4T_2$ can be prepared using the process and intermediates of the invention. The synthesis of this compound is shown in the Figures 1, 2 and 3 of this specification. As is described in the "Abstract" section of Wyatt, et al., *Proc. Natl. Acad. Sci.*, 1994, 91, 1356-1360, "The phosphorothioate oligonucleotide $T_2G_4T_2$ was identified as an inhibitor of HIV infection *in vitro*..."


Other preferred compounds having complementary sequences for herpes, papilloma and other viruses can also be prepared utilizing the process of the invention.

Oligonucleotides are also used as research reagents. For example, phosphorothioate oligonucleotides have been utilized in the studies of enzyme biochemistry and protein-nucleic acid interactions. These and other applications of phosphorothioate oligonucleotides are listed in "Oligonucleotides and Analogues A Practical Approach", F. Eckstein Ed., IRL Press, at pages 88-91, and


- 21 -

in the references contained therein, which are hereby incorporated by reference.

In one aspect, the present invention is directed to synthetic methods wherein a protected nucleoside having formula I is contacted with a coupling reagent such as one having the formula $(R_1R_2N)_2PO(CH_2)_xSiR_3R_4R_5$ for a time and under conditions effective to form a silylalkyl phosphoramidite monomer having formula II.



I



II

10

Such contacting preferably is effected under anhydrous conditions in the presence of a weak acid like 1H-tetrazole or 5-(4-nitrophenyl)-1H-tetrazole or diisopropylammonium tetrazolide or any other acid.

Q can be S, CH₂, CHF, CF₂ or, preferably, O. In a further preferred group of compounds, Q is S (see, e.g., Secrist, et al., Abstract 21, Synthesis and Biological Activity of 4'-Thionucleosides, Program & Abstracts, Tenth International Roundtable, Nucleosides, Nucleotides and their Biological Applications, Park City, Utah, Sept. 16-20, 1992). Each Q is independently selected and, hence, can be the same as or different from other Q within a given compound.

B_x can be nucleosidic bases selected from the natural nucleosidic bases adenine, guanine, uracil, thymine, cytosine as well as other bases including 2-aminoadenosine or 5-methylcytosine. In addition, other non-naturally occurring species can be employed to provide stable duplex or triplex formation with, for example, DNA. Representative

30

- 22 -

bases are disclosed in U.S. Patent No. 3,687,808 (Merigan et al.) which is incorporated herein by reference.

X can be OH, SH, SCH₃, F, OCN, O(CH₂)_nNH₂, O(CH₂)_nCH₃, where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, 5 substituted lower alkyl, alkaryl or aralkyl; Cl, Br, CN, CF₃, OCF₃, O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an 10 intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide. It is intended that the term "alkyl" denote branched and straight chain hydrocarbon residues, including alkyl groups having 15 one or more ³H and/or ¹⁴C atoms. It is preferred that X is H or OH, or, alternatively F, O-alkyl or O-alkenyl, especially where Q is O. Preferred alkyl and alkenyl groups have from 1 to about 10 carbon atoms.

Representative 2'-O-substituted oligonucleotides 20 are described in PCT patent application WO 93/13121, filed December 23, 1992, entitled Gapped 2' Modified Phosphorothioate Oligonucleotides, and U.S. patent application serial no. US93/06807 filed July 20, 1993 entitled Novel 2'-O-Alkyl Nucleosides and Phosphoramidites 25 Processes for the Preparation and Uses Thereof and in the references within these applications. Both applications are assigned to the same assignee as this application, and are hereby incorporated by reference.

For example, in the above cited references 2'-O- 30 substituted nucleosides are disclosed, including 2'-O -alkyl (methyl through octadecyl); -allyl, -dimethylallyl, -(N-phthalimido)prop-3-yl and -(N-phthalimido) pent-3-yl; -(imidazol-1-yl)butyl; and -pentyl-w-(N-phthalimido). Other substituent groups disclosed in the references as useful 35 for substitution at the 2' position of nucleosides include fluoro, C1-C9 aminoalkoxy; C1-C9 alkylimidazole, and polyethyleneglycol.

- 23 -

Y can be any hydroxyl protecting group for the 5' hydroxyl function of nucleosides. Preferably, Y is stable under basic conditions but can be removed under acidic conditions. A wide variety of protecting groups can be employed in the methods of the invention. In general, protecting groups render chemical functionality inert to specific reaction conditions, and can be appended to and removed from such functionality in a molecule. Representative protecting groups are disclosed by Beaucage, S.L.; Uyer, R.P., *Advances in the Synthesis of Oligonucleotides by the Phosphoramidite Approach*, Tetrahedron, 1992, 48, 2223-2311. Preferred protecting groups include dimethoxytrityl (DMTr), monomethoxytrityl, 9-phenylxanthen-9-yl (Pixyl) and 9-(p-methoxyphenyl)xanthen-9-yl (Mox).


Coupling reagents preferably have the formula $(R_1R_2N)_2PO(CH_2)_xSiR_3R_4R_5$ where R_1 and R_2 independently are alkyl having 1 to about 10 carbon atoms, R_3 , R_4 , and R_5 are, independently, alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbons atoms, and x is 1 to about 7. These are preferably prepared by reacting an alcohol having formula $HO(CH_2)_xSiR_3R_4R_5$ with phosphorus trichloride and reacting the resultant product, $Cl_2PO(CH_2)_xSiR_3R_4R_5$, with at least two equivalents of an amine having formula R_1R_2NH . The ammoniacal R groups can be the same or different and may be alkyl having 1 to about 10 carbon atoms, preferably 1 to 6 carbon atoms, more preferably 3 carbon atoms. Isopropyl groups are particularly preferred. The silyl R groups can be the same or different and may be alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbon atoms. Preferably, R_3 , R_4 and R_5 are selected from ethyl, ethyl, isopropyl, propenyl, n-butyl, t-butyl, and phenyl groups. Preferably, two of these silyl R groups are phenyl groups and one is a methyl group.

In the formula of the above coupling groups, the variable x can be 1 to about 7, preferably 1 to about 4, more preferably 2. A number of suitable alcohols are


- 24 -

disclosed by U.S. Patent No. 5,159,095, issued October 27, 1992 in the name of Celebuski, which is incorporated herein by reference. One preferred coupling reagent is diphenylmethyldisilylethyl N,N-diisopropylphosphoramidite, which can be derived from diphenylmethyldisilylethyl alcohol via diphenylmethyldisilylethyl phosphodichloridite.

Silylalkyl phosphoramidite monomers having formula II can be contacted with 3'-hydroxyl protected nucleosides having formula III for a time and under conditions effective to form phosphite dimers having formula IV, where the group W is a 3' hydroxyl protecting group.



III



IV

In preferred embodiments, such contact is effected under anhydrous conditions in the presence of an activating agent such as 1H-tetrazole, 5-(4-nitrophenyl)-1H-tetrazole, or diisopropylammonium tetrazolide.


W can be any hydroxyl protecting group for the 3' hydroxyl function of nucleosides. Preferably, W is stable under acidic conditions but can be removed under basic conditions. A wide variety of protecting groups can be employed in the methods of the invention. Preferred protecting groups include acyl protecting groups. A particularly preferred acyl group is acetyl. Other useful blocking groups that are stable under acid conditions but that can otherwise be removed are as is described in Beaucage et al., *ibid*.

- 25 -

Protecting groups for the compounds of the invention are selected such that the 3'-hydroxyl terminus protecting group W and the 5'-hydroxyl terminus protecting group Y are removed under different conditions. Protecting groups for the phosphite moiety, for example the $O(CH_2)_xSiR_3R_4R_5$ group of compounds of formula II, are chosen such that they are stable to conditions required for the removal of either the 3'-hydroxyl terminus protecting group W or the 5'-hydroxyl terminus protecting group Y.

Protecting groups for nucleobases are chosen such that they are stable to conditions required for the removal of either 3'-hydroxyl terminus protecting group W or 5'-hydroxyl terminus protecting group Y.

Phosphite compounds having formula IV can then be oxidized to produce compounds having formula V ($n = 1$).



Such oxidation can be effected to both phosphodiester ($Z = O$) and phosphorothioate ($Z = S$) structures. Useful sulfurizing agents include Beaucage reagent described in e.g., Iyer, R.P.; Egan, W.; Regan, J.B.; Beaucage, S.L., 3H-1,2-Benzodithiole-3-one 1,1-Dioxide as an Improved Sulfurizing Reagent in the Solid-Phase Synthesis of Oligodeoxyribonucleoside Phosphorothioates, *Journal of American Chemical Society*, 1990, 112, 1253-1254 and Iyer, R.P.; Phillips, L.R.; Egan, W.; Regan J.B.; Beaucage, S.L., The Automated Synthesis of Sulfur-Containing Oligodeoxyribonucleotides Using 3H-,2-Benzodithiol-3-one


- 26 -

- 1,1-Dioxide as a Sulfur-Transfer Reagent, *Journal of Organic Chemistry*, 1990, 55, 4693-4699. Tetraethyl- thiuram disulfide can also be used as described in Vu, H.; Hirschbein, B.L., Internucleotide Phosphite Sulfurization
- 5 With Tetraethylthiuram Disulfide, Phosphorothioate Oligonucleotide Synthesis Via Phosphoramidite Chemistry, *Tetrahedron Letters*, 1991, 32, 3005-3007. Further useful reagents for this step are dibenzoyl Tetrasulfide, Rao, M.V.; Reese, C.B.; Zhengyun, Z., Dibenzoyl Tetrasulphide - A
- 10 Rapid Sulphur Transfer Agent in the Synthesis of Phosphorothioate Analogues of Oligonucleotides, *Tetrahedron Letters*, 1992, 33, 4839-4842; di(phenylacetyl)disulfide, Kamer, R.C.R.; Roelen, H.C.P.F.; van den Eist, H.; van der Marel, G.A.; van Boom, J.H., An Efficient Approach Toward
- 15 the Synthesis of Phosphorothioate Diesters Via the Schonberg Reaction, *Tetrahedron Letters*, 1989, 30, 6757-6760; sulfur; and sulfur in combination with ligands like triaryl, trialkyl or triaralkyl or trialkaryl phosphines. Useful oxidizing agents include
- 20 iodine/tetrahydrofuran/water/pyridine, hydrogen peroxide/water, tert-butyl hydroperoxide, or a peracid like m-chloroperbenzoic acid. In the case of sulfurization, reaction is performed under anhydrous conditions with the exclusion of air, in particular oxygen, whereas in the case
- 25 of oxidation the reaction can be performed under aqueous conditions.

In certain embodiments, compounds having formula V are exposed to reaction conditions effective to remove the 5' hydroxyl protecting group Y, and the resultant product is

30 contacted with additional monomer II to form phosphite oligonucleotides having formula VIII ($n = 1$).


- 27 -



As will be recognized, further compounds having
 5 formula VIII, wherein Y and W are hydroxyl protecting groups
 and n is, for example, 2-200, can be prepared by oxidizing
 the phosphite intermediate to the phosphate, followed by
 removing the Y hydroxyl protecting group, and reacting with
 additional monomer II. It will further be recognized, that
 10 multiple iteration of this reaction can be effect to extend
 the oligonucleotide for increasing values of n.

In preferred embodiments, compound having formula
 IV are exposed to reaction conditions effective to remove
 the 5' hydroxyl protecting group Y, and reacting the
 15 resultant product is with additional monomer II to form
 phosphite oligonucleotides having formula IVb (n = 2).


- 28 -



IVb

As will be recognized, further compounds having
 5 formula IVb, wherein W is a hydroxyl protecting group and n
 is, for example, 3-200, can be prepared by removing the 5'-
 hydroxyl protecting group Y and adding additional monomer of
 formula II.

Oxidation products having, for example, formula V
 10 can be exposed to conditions effective to cleave the 3'-OH
 protecting group W. The resulting product is contacted with
 ammonium hydroxide or some other aqueous base or fluoride
 ion to remove the silylalkyl portion thereof to yield
 phosphodiester and phosphorothioate-containing compounds
 15 having, for example, formula IX.



IX

- 29 -

Contact with fluoride ion preferably is effected in a solvent such as tetrahydrofuran or acetonitrile or dimethoxyethane or water. Fluoride ion preferably is provided in the form of one or more salts selected from

5 tetraalkylammonium fluorides (e.g., tetrabutylammonium fluoride (TBAF) or potassium fluoride or Cesium fluoride.

A more preferred method of removing the phosphorous protecting group is effected via the use of tetrafluorosilane in a "wet" solvent. Suitable as the

10 solvent is acetonitrile having a trace of water. A 100:1 mixture gave essentially immediate deprotection at room temperature without any indication of cleavage of the internucleotide linkage. While we do not wish to be bound by theory, it is believed that this deprotection is effected

15 via a β -fragmentation mechanism that is a result of coordination of the silicon atom of the tetrafluorosilane with the ether oxygen atom of the protecting group. Electron withdrawal at the ether oxygen atom is augmented by the electron withdrawing effects of the fluorine atoms of

20 the tetrafluorosilane moiety.

The 3'-hydroxyl protecting group Y can be removed from compounds having, for example, formula IX by techniques well known in the art to produce compounds having formulas VI and VIa wherein Y is replaced by H. For example,

25 dimethoxytrityl protecting groups can be removed by protic acids such as formic acid, dichloroacetic acid, trichloroacetic acid, p-toluene sulphonic acid or any other acid or with Lewis acids like zinc bromide or any other Lewis acid.

30 The 3'-hydroxyl protecting group W can be removed from compounds having, for example, formula IX by techniques well known in the art to produce compounds wherein W is replaced by H. For example, acetyl protecting groups can be removed by barium hydroxide.

35 In a preferred embodiment, compounds of the present invention are reacted in iterative fashion to produce oligonucleotides of up to about 200 units. A more

- 30 -

preferred size is up to about 50 units. Figure 1 is a reaction schematic depicting exemplary routes for the preparation of intermediates useful in the synthesis of oligonucleotides. 3'-O-Acetyl deoxythymidine is reacted with 5'-DMT DPSE phosphoramidite, monomer 2, and subsequently oxidized to form dimer 3, which has hydroxyl protecting groups at the 3'- and 5'-hydroxyl termini. Dimer 3 is then either deprotected at the 5'-hydroxyl to yield dimer 4, or deprotected at the 3'-hydroxyl and phosphitylated to produce activated dimer 5. 3'-O-Acetyl deoxyguanine is reacted with 5'-DMT DPSE phosphoramidite 6 and subsequently sulfurized, i.e. oxidized, to form dimer 7, which is then either deprotected at the 5'-hydroxyl to yield dimer 8, or deprotected at the 3'-hydroxyl and phosphitylated to produce activated dimer 9.

Figure 2 shows the coupling of dimer 8 and activated dimer 5 and subsequent sulfurization, 3'-O-deacetylation, and phosphitylation to yield activated tetramer 10. Figure 2 also shows the coupling of dimer 4 and activated dimer 9 and subsequent sulfurization and 5'-hydroxyl deprotection to yield 3'-O-acetylated tetramer 11.

Figure 3 shows the coupling and subsequent sulfurization of 3'-O-acetylated tetramer 11 and activated tetramer 10 to yield the fully protected octamer 12, and subsequent reaction removal all protecting groups to yield oligomer T-T-G-G-G-G-T-T. As will be recognized, oligomers containing odd numbers of nucleotide units may be synthesized by incorporating a monomer of formula II or another nucleosidic moiety containing an odd number of nucleoside subunits into the iterative process at any point in the process.

It will also be recognized that the processes of the invention may also be accomplished by reaction of a nucleotide containing a free hydroxyl at the 3' terminus with a nucleosidic moiety phosphitylated at the 5'-hydroxy terminus.

- 31 -

Additionally, it will be recognized that the process steps of the present invention need not be performed any particular number of times or in any particular sequence. Accordingly, products incorporating one or
5 several functional groupings of the enumerated formulas may be constructed according to the invention. For example, the iterative oxidation of a growing oligonucleotide chain may be independently achieved with sulfurizing agents or oxidizing agents to yield oligomers containing both
10 phosphate and phosphorothioate groups.

It will be further seen that by substitution of the alkylsilyl ethoxy group of the coupling reagent $(R_1R_2N)_2PO(CH_2)_xSiR_3R_4R_5$ the process of the present invention may be employed to make other phosphorous-substituted
15 oligonucleotides. For example, substitution of one or more of the alkylsilylethoxy groups with methyl groups would result in a methylphosphite intermediate, capable of subsequent oxidation by the methods of the invention to yield an oligonucleotide containing one or more methyl
20 phosphonate linkages. In a like manner, use of β -cyanoethyl phosphorous groups can be employed. Analogously, it may be seen that starting with a phosphorothioamidite, the process of the invention may be utilized to produce oligonucleotides containing one or more phosphorodithioate linkages.

25 Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following example thereof, which are not intended to be limiting.

5'-O-(4,4'-Dimethoxytrityl)-3'-O-[(N,N-
30 diisopropylamino)-2-(diphenylmethyl-silyl)ethoxyphosphino]-thymidine was synthesized according to a published procedure: see V. T. Ravikumar, T. K. Wyrzykiewicz, D. L. Cole, *Tetrahedron* **1994**, 50(31), 9255-9266. 5'-O-(4,4'-
Dimethoxytrityl)-3'-O-[(N,N-diisopropylamino)-2-(
35 diphenylmethyl-silyl)ethoxyphosphino]-deoxyguanosine was synthesized according to a published procedure: see V. T. Ravikumar, T. K. Wyrzykiewicz, D. L. Cole, *Tetrahedron* **1994**,

- 32 -

50(31), 9255-9266. 3'-O-Acetyl-thymidine was purchased from Sigma. 3'-O-Levunilyl-thymidine was synthesized according a published procedure: see G. Kumar, M. S. Poonian, *J. Org. Chem.* 1984, 49, 4905-4912.

5 EXAMPLE 1

Preparation of Fully Protected TT Dimer:

To a solution of 1.42 g (5.0 mmol) of 3'-acetyl thymidine and 0.35 g (4.0 mmol) of 1-H Tetrazole in acetonitrile (60 mL) is added 5.50 g (6.0 mmol) of 5'-O-
10 (4,4'-dimethoxytrityl)thymidine-3'-O-(2-diphenylmethylsilylethyl N,N-diisopropylphosphoramidite in 40 mL acetonitrile. The reaction mixture is stirred at room temperature under argon for 0.5 h. A solution of 5.0 g (25 mmol) of 3H-1,2-benzodithiol-3-one 1,1-dioxide in
15 acetonitrile is added as quickly as possible with vigorous stirring. The reaction mixture is stirred at room temperature for 20 minutes. The reaction mixture is then filtered and concentrated. The crude product is purified by flash chromatography on silica gel using ethyl
20 acetate/hexane. 1% triethylamine is used during the purification. ³¹P-NMR (CDCl₃, ppm): 66.80, 67.07.

EXAMPLE 2

Preparation of Fully Protected GG dimer:

To a solution of 3'-acetyl-N²-isobutyryl-2'-
25 deoxyguanosine (5.0 mmole) and 1-H Tetrazole (4.0 mmole) in acetonitrile (60 mL) is added 5.50 g (6.0 mmol) of N²-Iso-butyryl-5'-O-(4,4'-dimethoxytrityl)-2'-deoxyguanosine-3'-O-(2-diphenylmethylsilylethyl N,N-diisopropylphosphoramidite) in acetonitrile (40 mL). The reaction mixture is stirred at
30 room temperature under argon for 0.5 h. A solution of 5.0 g (25 mmole) of 3H-1,2-benzodithiol-3-one 1,1-dioxide in acetonitrile is added as quickly as possible with vigorous stirring. The reaction mixture is stirred at room temperature for 20 minutes. The reaction mixture is then
35 filtered and concentrated. The crude product is purified by

- 33 -

flash chromatography on silica gel using ethyl acetate/hexane. 1% Triethylamine is used during the purification. ^{31}P -NMR (CDCl_3 , ppm): 66.8, 67.1.

EXAMPLE 3**5 Preparation of the 5'-HO-TT Dimer:**

Fully protected phosphorothioate TT dimer (1 mmol) is dissolved in dichloromethane (50 mL) and 3% dichloroacetic acid in dichloromethane (v/v) (20 mL) is added and stirred for 15 minutes. The reaction mixture is concentrated and purified by flash chromatography on silica gel using ethyl acetate/hexane. 1% Triethylamine is used during the purification. ^{31}P -NMR (CDCl_3 , ppm): 67.3, 67.41.

EXAMPLE 4**Preparation of the 5'-HO-GG Dimer:**

15 Fully protected phosphorothioate GG dimer (1 mmol) is dissolved in dichloromethane (50 mL) and 3% dichloroacetic acid in dichloromethane (v/v) (20 mL) is added and stirred for 15 minutes. The reaction mixture is concentrated and purified by flash chromatography on silica gel using ethyl acetate/hexane. 1% Triethylamine is used during the purification. ^{31}P -NMR (CDCl_3 , ppm): 67.5, 67.6.

EXAMPLE 5**Preparation of Fully Protected TTT trimer:**

To a stirred solution of 5'-HO-TT phosphorothioate dimer (5.0 mmole) and 1-H Tetrazole (5.0 mmole) in acetonitrile (60 mL) is added 5'-O-(4,4'-dimethoxytrityl) thymidine-3'-O(2-diphenylmethylsilylethyl N,N-diisopropylphosphoramidite) (6.0 mmol) in acetonitrile (40 mL). The reaction mixture is stirred at room temperature under argon for 0.5 h. A solution of 3H-1,2-benzodithiol-3-one 1,1-dioxide (25 mmol) in acetonitrile is added as quickly as possible with vigorous stirring. The reaction mixture is stirred at room temperature for 20 minutes. The reaction mixture is then filtered and concentrated. The crude

- 34 -

product is purified by flash chromatography on silica gel using ethyl acetate/hexane. 1% Triethylamine is used during the purification.

EXAMPLE 6**5 Preparation of Fully Protected CTT trimer:**

To a solution of 5'-HO-TT phosphorothioate dimer (5.0 mmole) and 1-H Tetrazole (5.0 mmole) in acetonitrile (60 mL) is added N⁴-Benzoyl-5'-O-(4,4'-dimethoxytrityl)-2'-deoxycytidine-3'-O-(2-diphenylmethylsilylethyl N,N-diisopropylphosphoramidite) (6.0 mmol) in acetonitrile (40 mL). The reaction mixture is stirred at room temperature under argon for 0.5 h. A solution of 3H-1,2-benzodithiol-3-one 1,1-dioxide (25 mmol) in acetonitrile is added as quickly as possible with vigorous stirring. The reaction mixture is stirred at room temperature for 20 minutes. The reaction mixture is then filtered and concentrated. The crude product is purified by flash chromatography on silica gel using ethyl acetate/hexane. 1% Triethylamine is used during the purification.

20 EXAMPLE 7**5'-O-(4,4'-Dimethoxytrityl)-3'-(O-acetyl)-thymidine dimer**

A mixture of 5'-O-(4,4'-dimethoxytrityl)-3'-O-[(N,N-diisopropylamino)-2-(diphenylmethylsilyl)ethoxyphosphino]-thymidine (1.83 g, 2.0 mmol), 3'-(O-acetyl)-thymidine (284 mg, 1.0 mmol) and 1H-tetrazole (560 mg, 8.0 mmol) was dried under vacuum and exposed to inert gas (argon). Dry acetonitrile (8 mL) was added. After 3 h at room temperature, 3H-1,2-benzodithiol-3-one-1,1-dioxide (1.0 g, 5.0 mmol) was added. After 30 min, triethylamine (1 mL) was added to the mixture and the mixture was filtered. The filtrate was concentrated under reduced pressure and the oily residue was purified by column chromatography (silica, 1. ethyl acetate/n-hexanes 1:1, then 8:2) to give the title compound (1.0 g, 88%). ³¹P NMR (CDCl₃): 67.4, 67.5 ppm.

- 35 -

EXAMPLE 8**5'-(O-4,4'-Dimethoxytrityl)-3'-(O-levunilyl)-thymidine dimer**

A mixture of 5'-O-(4,4'-dimethoxytrityl)-3'-O-
[(*N,N*-diisopropylamino)-2-(diphenylmethyl-silyl)ethoxy-
5 phosphino]-thymidine (31.73 g, 34.6 mmol), 3'-(O-levunilyl)-
thymidine (10.21 g, 30.0 mmol) and 1*H*-tetrazole (8.4 g, 120
mmol) was dried under vacuum and exposed to inert gas
(argon). Dry acetonitrile (150 ml) was added. After 4 h at
room temperature, 3*H*-1,2-benzodithiol-3-one-1,1-dioxide
10 (20.0 g, 100.0 mmol) was added. After 2h, triethylamine (20
ml) was added and the mixture was filtered. The filtrate was
concentrated under reduced pressure and the oily residue
purified by column chromatography (silica, ethyl acetate) to
give the title compound (37 g). ³¹P NMR (CDCl₃): 67.4, 67.6
15 ppm.

EXAMPLE 9**5'-(O-4,4'-Dimethoxytrityl)-thymidine dimer**

5'-(O-4,4'-Dimethoxytrityl)-3'-(O-levunilyl)-
thymidine dimer (35.0 g) was dissolved in an ice-cold
20 solution of hydrazine-hydrate (10.0 g), pyridine (240 ml)
and acetic acid (240 ml). After 8 min, ice was added,
followed by extraction with dichloromethane. The organic
phase was kept over Na₂SO₄, filtered and the solvent was
removed. The residue was purified by column chromatography
25 (silica, 1. ethyl acetate/*n*-hexanes 1:1, then ethyl acetate,
0.1% triethylamine) to give the title compound (24.9 g, 81
%). ³¹P NMR (CDCl₃): 66.8, 67.7 ppm.

EXAMPLE 10**5'-(O-4,4'-Dimethoxytrityl)-thymidine dimer amidite**

30 Under argon, a solution of 1*H* tetrazole (35 mg,
0.5 mmol) and 2-cyanoethyl-*N,N,N',N'*-tetra-
isopropylphosphorodiamidite (3.0 mmol) in dry acetonitrile
(10 ml) was added to thymidine dimer of Example 9 (2.18 g,
2.0 mmol). After amidite formation was complete,
35 triethylamine (0.5 ml) and ethyl acetate were added. The

- 36 -

solution was extracted with aqu. NaHCO_3 (2%). The organic phase was kept over Na_2SO_4 and the solvent was removed under reduced pressure. The residue was purified by column chromatography (silica, ethyl acetate, 1% triethylamine) to give the phosphoramidite title compound (2.2 g, 85 %). ^{31}P NMR (CDCl_3): 67.66, 67.73, 67.84, 149.47, 149.68, 149.98 ppm.

EXAMPLE 11**3'-(O-levunilyl)-thymidine dimer**

The 5'-(O-4,4'-dimethoxytrityl)-3'-(O-levunilyl)-thymidine dimer from Example 8 (30.5 g) was dissolved in a solution of dichloroacetic acid (4.3 ml) and dichloromethane (80 ml). After 10 min, ethanol (6.0 g) and triethyl silane (12.0 g) were added. After the orange color had disappeared, the solution was extracted with water and NaHCO_3 (1M) solution. The organic layer was kept over Na_2SO_4 . After filtration, the solvent was removed and the residue was purified by column chromatography (silica, 1. ethyl acetate/n-hexanes 1:1, then ethyl acetate) to give the title compound (20.7 g). ^{31}P NMR (CDCl_3): 66.82, 67.67 ppm.

EXAMPLE 12**5'-(O-4,4'-Dimethoxytrityl)-3'-(O-levunilyl)-dG-T-T- trimer**

Under argon, a mixture of 1H-tetrazole (350 mg, 5.0 mmol) and dG amidite, i.e. 5'-O-(4,4'-dimethoxytrityl)-3'-O-[(N,N-diisopropylamino)-2-(diphenylmethyl-silyl)-ethoxyphosphino]-deoxyguanosine, was dissolved in dry acetonitrile (5 ml). After 15 min, a solution of thymidine dimer from Example 11 (885 mg, 1.0 mmol) in dry acetonitrile (2.5 ml) was added. After 15 min, the solution was cooled to 0 °C and a solution of 3H-1,2-benzodithiol-3-one-1,1-dioxide (1.2 g, 6.0 mmol) were added. After 15 min, triethylamine (0.5 ml) was added and the mixture was filtered. The filtrate was concentrated under reduced pressure and the oily residue was dissolved in ethyl acetate and extracted with aqu. NaHCO_3 (1M). The organic phase was kept over Na_2SO_4 , filtered and the filtrate was concentrated under reduced

- 37 -

pressure. The residue was purified by column chromatography (silica, ethyl acetate) to give the title compound (1.59 g). ^{31}P NMR (CDCl_3): 66.9, 67.0, 67.3, 68.0, 68.3 ppm.

EXAMPLE 135 **5'-(O-4,4'-Dimethoxytrityl)-dG-T-T- trimer**

5'-(O-4,4'-Dimethoxytrityl)-3'-(O-levunilyl)-dG-T-T- trimer from Example 12 (1.43 g, 0.783 mmol) was dissolved in ice-cold hydrazine hydrate (0.3 ml), pyridine (7.2 ml) and acetic acid (4.8 ml). After 1 h, ice (65 g) was added.
10 The title trimer precipitated and was isolated by filtration and rinsed with water, yield 1.3 g.

EXAMPLE 14**3'-O-Levulinyl-N2-isobutyryl-2'-deoxyguanosine**

5'-DMT-N2-isobutyryl-2'-deoxyguanosine (71.6 g, 0.112 mol) was transferred to a 1000 ml flask and anhydrous dioxane (700 ml) was added and stirred until the solution became homogeneous. Dicyclohexylcarbodiimide (57.8 g, 0.280 mol), levulinic acid (25.9 g, 0.224 mol) and 4-dimethylaminopyridine (0.56 g) were added and the mixture
20 vigorously stirred using magnetic stirring. After 3 h, the reaction mixture was filtered, the solid residue washed with ethyl acetate (250 ml). The filterates were combined and concentrated to afford a product.

This product was dissolved in dichloromethane (400 ml) and 2.5% DCA in dichloromethane (160 ml) was added and stirred. After 1 h, the reaction mixture was diluted with dichloromethane (400 ml) and washed with saturated sodium bicarbonate solution. The organic layer was separated, dried and concentrated. The crude product was purified by
30 flash chromatography using silica gel to give the title compound as a colorless solid, 35.1 g (80%).

- 38 -

EXAMPLE 15**Synthesis of 5'-HO-GG-OLev.**

5'-O-(4,4'-dimethoxytrityl)-N2-isobutyryl-2'-deoxyguanosine-3'-O-(2-diphenylmethylsilylethyl) N,N-diisopropyl phosphoramidite (36 g, 0.0356 mol) and 1H-tetrazole (8 g, 0.114 mol) were charged into a 1000 ml flask provided with magnetic stirring. Anhydrous acetonitrile (350 ml) was added and the mixture stirred. To this homogenous solution, 3'-O-levulinyl-N2-isobutyryl-2'-deoxyguanosine in acetonitrile (200 ml) was added with stirring. After 2 h, a solution of Beaucage reagent (28.5 g, 0.142 mol) in acetonitrile (90 ml) was added as quickly as possible with vigorous stirring. After 1 h, the reaction mixture was filtered and concentrated. This crude product was redissolved in dichloromethane (300 ml) and a solution of 2.5% DCA in dichloromethane was added and stirred. After 1 h, the reaction mixture was washed with saturated sodium bicarbonate solution, dried and concentrated. The crude product was purified by flash chromatography using silica gel to give the title compound as a colorless solid.

EXAMPLE 16**Synthesis of 5'-HO-TGG-OLev.**

5'-O-(4,4'-dimethoxytrityl)thymidine-3'-O-(2-diphenylmethylsilylethyl) N,N-diisopropyl phosphoramidite (0.02488 mol) and 1H-tetrazole (0.1 mol) were charged into a 1000 ml flask provided with magnetic stirring. Anhydrous acetonitrile (350 ml) was added and the mixture stirred. To this homogenous solution, 5'-HO-GG-OLev (0.02 mol) in acetonitrile (200 ml) was added with stirring. After 2 h, a solution of Beaucage reagent (0.1 mol) in acetonitrile (90 ml) was added as quickly as possible with vigorous stirring. After 1 h, the reaction mixture was filtered and concentrated. This crude product was redissolved in dichloromethane (300 ml) and a solution of 2.5% DCA in dichloromethane was added and stirred. After 1 h, the reaction mixture was washed with saturated sodium

- 39 -

bicarbonate solution, dried and concentrated. The crude product was purified by flash chromatography using silica gel to give the title compound as a colorless solid.

EXAMPLE 17**5 Synthesis of 5'-HO-CTGG-OLev.**

5'-O-(4,4'-dimethoxytrityl)-N⁴-acetyl-2'-deoxycytidine-3'-O-(2-diphenylmethylsilylethyl) N,N-diisopropyl phosphoramidite (0.02488 mol) and 1H-tetrazole (0.1 mol) are charged into a 1000 ml flask provided with magnetic stirring. Anhydrous acetonitrile (350 ml) is added with stirring. To this homogenous solution, 5'-HO-TGG-OLev (0.02 mol) in acetonitrile (200 ml) is added and stirred. After 2 h, a solution of Beaucage reagent (0.1 mol) in acetonitrile (90 ml) is added as quickly as possible with vigorous stirring. After 1 h, the reaction mixture is filtered and concentrated. This crude product is redissolved in dichloromethane (300 ml) and a solution of 2.5% DCA in dichloromethane is added and stirred. After 1 h, the reaction mixture is washed with saturated sodium bicarbonate solution, dried and concentrated. The crude product is purified by flash chromatography using silica gel to give the title compound.

EXAMPLE 18**Synthesis of 5'-DMT-GCTGG-OLev.**

5'-O-(4,4'-dimethoxytrityl)-N²-isobutyryl-2'-deoxyguanosine-3'-O-(2-diphenylmethylsilylethyl) N,N-diisopropyl phosphoramidite (0.02488 mol) and 1H-tetrazole (0.1 mol) are charged into a 1000 ml flask provided with magnetic stirring. Anhydrous acetonitrile (350 ml) is added with stirring. To this homogenous solution, 5'-HO-CTGG-OLev (0.02 mol) in acetonitrile (200 ml) is added and the mixture is stirred. After 2 h, a solution of Beaucage reagent (0.1 mol) in acetonitrile (90 ml) is added as quickly as possible with vigorous stirring. After 1 h, the reaction mixture is filtered and concentrated. The crude

- 40 -

product is purified by flash chromatography using silica gel to give the title compound.

EXAMPLE 19

Use of repeating blocks in the preparation of multiple 5 oligonucleotides

In a further aspect of the invention methods are provided for the preparation of a plurality of oligonucleotides (target oligonucleotides) which contain common nucleotide subsequences. According to the method,
10 the two or more target oligonucleotides to be synthesized are analyzed to identify occurrences of common nucleotide subsequences that occur in each target oligonucleotide. The subsequences can be dimers, trimers, tetramers, pentamers, or any such "short mer."

15 For example, in the target oligonucleotides:

GCT GGT GGC CTG GA SEQ ID No. 1

and

CTG GTG GGC TGG TGG A SEQ ID No. 2

the trimer TGG occurs in each of the target oligonucleotides
20 (once in sequence No.1 and twice in sequence No. 2), the tetramer CTGG occurs once in each of sequence No. 1 and No. 2, as does the pentamer GCTGG. Conveniently, the nucleotides can be regrouped, and the subsequences can be visualized as:

25 GCTGG TGG C CTGG A for SEQ No. 1; and

CTGG TGG GCTGG TGG A for SEQ No. 2.

thus facilitating visualization of the various subsequences within the target oligonucleotides.

The analysis can be effected by visualization, or
30 appropriate software routines can be used to select the subsequences. Such analyses can include certain statistical analysis regimes to select the greatest occurrence of the least number of subsequences, or other factors such as the least number of machine cycles required for synthesis. Thus
35 for SEQ No. 2 an alternate pattern could have been selected, such as, for example:


- 41 -

C TGG TGG GC TGG TGG A for SEQ No. 2.

This second pattern for this same sequence would allow for repeated use of the trimer block "TGG" four times. However, a total of six machine coupling cycles would have to be effected using this strategy. Because it would yield the least number of machine cycles, and further includes subsequences that correspond to subsequences in SEQ No. 1, the first analysis pattern for SEQ No. 2 is selected for further illustration in this example.

Using the methods of Examples 1-18, synthons the following subsequences are prepared: a trimer block corresponding to the nucleotide subsequence TGG; a tetramer block corresponding to the nucleotide subsequence CTGG; and a pentamer block corresponding to the nucleotide subsequence GCTGG.

The trimer, tetramer and pentamer oligonucleotide subsequences are each phosphitylated according to the procedures of the above examples to form oligonucleotide synthons having the general formula:



wherein the constituent members are as defined above.

According to the method, the oligonucleotide synthons are reacted with nucleoside precursors to effect the synthesis of target oligonucleotides. Nucleoside precursors, as used herein, are standard reagents used for solid phase or solution phase nucleotide synthetic

- 42 -

methodologies, including phosphotriester, phosphite triester, and H-phosphonate chemistries. A wide variety of nucleotide precursors, such as, for example, nucleoside phosphoramidites, are commercially available from suppliers
5 such as Sigma Chemical Co., St. Louis Mo.

To synthesize the target oligonucleotide of SEQ No. 1 a standard commercial oligonucleotide CPG solid support having a first nucleotide A is utilized as the support. The A nucleotide residue on the support will form
10 the 3' terminus of the target oligonucleotide upon cleavage of the completed target oligonucleotide from the support. The target nucleotide is completed by adding first one unit of the CTGG tetramer block, followed by a unit of a standard C nucleotide phosphoramidite, followed by a unit of the TGG
15 trimer block and finally one unit of the GCTGG pentamer block. Thus, to form the 14 mer, only four phosphoramidite units are needed and only four machine cycles are required. To synthesize the target oligonucleotide of SEQ No. 2, a CPG solid support that includes a standard A nucleotide unit
20 thereon is selected for the support. To this is added a first unit of the TGG trimer block, followed by a unit of GCTGG pentamer block, next a second unit of the TGG trimer block and finally a unit of the CTGG tetramer block. Thus for this 16 mer target oligonucleotide four separate
25 phosphoramidite units and four machine cycles are needed to complete the synthesis.

The target oligonucleotides can be synthesized utilizing any of the standard backbones such as phosphodiester, phosphorothioate, phosphorodithioate, methyl
30 phosphonates and the like, as is standard for machine aided oligonucleotide synthesis, by choice of either the starting amidite or by selection of the appropriate oxidization agent. Those skilled in the art will appreciate that numerous changes and modifications may be made to the
35 preferred embodiments of the invention and that such changes and modifications may be made without departing from the spirit of the invention. It is therefore intended that the


- 43 -

appended claims cover all such equivalent variations as fall within the true spirit and scope of the invention.


- 44 -

What is claimed is:

1. A method for the solution phase preparation of an oligonucleotide comprising reacting, in solution, a first synthon having the structure




with a second synthon having the structure



to form a moiety having the structure

- 45 -



where each Q is independently O, S, CH₂, CHF or CF₂;
 each B_x is independently a nucleosidic base;
 each X is independently, OH, SH, SCH₃, F, OCN, O(CH₂)_nNH₂, O(CH₂)_nCH₃ where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl or aralkyl; Cl, Br, CN, CF₃, OCF₃, O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a conjugate; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide;

each Y is independently a 5' hydroxyl protecting group;

W is a 3' hydroxyl protecting group;

each Z is independently O or S;

each T is independently a phosphorous blocking group;

U is a phosphite activating group; and


n is an integer from 0 to 50.

- 46 -

2. The process of claim 1 wherein each group T is $R_3R_4R_5$ silylalkoxy wherein R_3 , R_4 and R_5 are alkyl or aryl.
3. The process of claim 2 wherein each R is, independently, methyl or phenyl.
4. The process of claim 1 wherein U is a dialkylamino group.
5. The process of claim 1 wherein said second synthon is formed by reacting a reagent $(R_1R_2N)_2PO(CH_2)_xSiR_3R_4R_5$, wherein R_1 and R_2 independently are alkyl having 1 to about 10 carbon atoms, and R_3 , R_4 , and R_5 are, independently, alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbons atoms, and x is 1 to about 7; with a nucleoside to form said second synthon.
6. The process of claim 5 wherein the reaction takes place in the presence of 1H-tetrazole, 5-(4-nitrophenyl)-1H-tetrazole, or diisopropylammonium tetrazolide.
7. The process of claim 1 further comprising removing the groups W, T, and Y from the moiety and oxidizing the moiety to form either phosphorothioate or phosphodiester bonds.
8. The process of claim 1 further comprising transforming the moiety into a first synthon for iterative reaction with a further second synthon.
9. The process of claim 1 further comprising transforming the moiety into a second synthon for iterative reaction with a further first synthon.
10. A method for preparing an oligomer comprising reacting together, in solution,

- 47 -

a first synthon comprising at least two nucleoside units and having a 5' location protected with a 5' hydroxylic blocking group and a 3' location substituted with a function having the formula



wherein U is a phosphite activating group and T is a phosphorous blocking group,

with a second synthon comprising a nucleoside unit having a 3' location protected with a 3' hydroxylic blocking group and a 5' location capable of reacting with the U-P-T function.



11. The method of claim 10 wherein the product of the reaction is oxidized to form either phosphate or phosphorothioate internucleoside bonds.

12. The method of claim 10 wherein said second synthon comprises at least two nucleoside units.

13. The method of claim 10 wherein the function U-P-T is incorporated through reaction with a reagent $(\text{R}_1\text{R}_2\text{N})_2\text{PO}(\text{CH}_2)_x\text{SiR}_3\text{R}_4\text{R}_5$ wherein R_1 and R_2 independently are alkyl having 1 to about 10 carbon atoms, R_3 , R_4 , and R_5 are, independently, alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbons atoms, and x is 1 to about 7.

- 48 -

14. A compound having one of the formulas



where each Q is independently O, S, CH₂, CHF or CF₂;
 each B_x is independently a nucleosidic base;
 each X is independently, OH, SH, SCH₃, F, OCN,
 O(CH₂)_nNH₂, O(CH₂)_nCH₃, where n is from 1 to about 10; C₁ to C₁₀
 lower alkyl, substituted lower alkyl, alkaryl or aralkyl;
 Cl, Br, CN, CF₃, OCF₃, O-, S-, or N-alkyl; O-, S-, or N-
 alkenyl; SOCH₃, SO₂CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl;
 heterocycloalkaryl; aminoalkylamino; polyalkylamino; sub-
 stituted silyl; an RNA cleaving group; a conjugate; a
 reporter group; an intercalator; a group for improving the
 pharmacokinetic properties of an oligonucleotide; or a group
 for improving the pharmacodynamic properties of an
 oligonucleotide;

each Z is independently O or S;

Y is H or a hydroxyl protecting group;

U is a phosphite activating group;

each T is independently a phosphorous blocking
 group; and

n is an integer from 1 to 200.

15. The compound of claim 14 wherein Q is O.

16. The compound of claim 14 wherein X is H, OH or O-alkyl.


- 49 -

17. The compound of claim 14 wherein T has the formula $-O(CH_2)_xSiR_3R_4R_5$ wherein R_3 , R_4 , and R_5 are, independently, alkyl having 1 to about 10 carbon atoms or aryl having 6 to about 10 carbons atoms, and x is 1 to about 7.

18. A library comprising a plurality of compounds in accordance with claim 14.

19. A method for the preparation of a plurality of target oligonucleotides, each of said target oligonucleotides comprising at least one common nucleotide subsequence, comprising:

identifying said common nucleotide subsequences;
providing an oligonucleotide synthon of formula:



wherein:

each Q is O;

each B_x is independently a nucleosidic base;
each X is independently, H, OH, F, O-alkyl or substituted O-alkyl wherein said substitution is amino, alkoxy or nitrogen heterocycle;

each Z is independently O or S;

Y is H or a hydroxyl protecting group;

U is a phosphite activating group;

- 50 -

each T is independently a phosphorous blocking group; and

n is an integer from 1 to 200;

said oligonucleotide synthon having one of said common nucleotide subsequences; and

reacting said synthon with nucleoside precursors to form said target oligonucleotide.

20. The process of claim 19 wherein a plurality of said synthons are so reacted.

FIG. 1







FIG.2

2/3



3/3

FIG.3



INTERNATIONAL SEARCH REPORT

Int. national application No.
PCT/US95/06825

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C07H 1/00, 19/06, 19/16

US CL : 536/ 23.1, 25.3, 25.34

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/ 23.1, 25.3, 25.34

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE

search terms: phosphoramidite, silylalkyl, oligonucleotide

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 5,113,005 (CELEBUSKI) 12 May 1992, columns, 1-3, column 9, lines 16-62, column 10, lines 6-51, column 11, lines 34-53.	1-20
Y	US, A, 4,458,066 (CARUTHERS ET AL.) 03 July 1984, column 6, line 25 through column 7, line 35, column 8, line 36 through column 9, line 22.	1-20

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>		<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>
--	--	---

Date of the actual completion of the international search

10 JULY 1995

Date of mailing of the international search report

18 AUG 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

JAMES O. WILSON

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*